

LAB TEST #2

Directions: Answer the following questions in the spaces provided. Show all work and include units on your final answers to receive full credit. Pay attention to significant digits!

1. (15 points) You wish to determine the heat capacity of a coffee cup calorimeter. You may do so by mixing hot and cold water in a polystyrene coffee-cup, measuring the temperature changes, and then applying the principles of calorimetry. The following data were collected:


Determination of the heat capacity of a calorimeter

| | |
|---|---------|
| Volume of cold water | 50.8 mL |
| Temperature of cold water <i>The cold water is initially in the calorimeter.</i> | 23.7°C |
| Volume of hot water | 49.6 mL |
| Temperature of hot water | 56.9°C |
| Final temperature of the mixture | 38.1°C |


The density of the water is 1.00 g/mL, and the specific heat of water is $4.18 \text{ J/g}\cdot\text{C}^\circ$.

What is the heat capacity of the calorimeter?

2. (18 points) You mix a bleach sample (NaOCl is the active ingredient) with potassium iodide and acid to completely form triiodide (I_3^-) ions by the following reaction:



You then titrate the I_3^- ions with a sodium thiosulfate ($Na_2S_2O_3$) solution you have previously determined the concentration of. The triiodide and the sodium thiosulfate react by the following equation:



The following data were collected:

Titration of Bleach sample

| | |
|--|----------|
| [$Na_2S_2O_3$] (determined previously) | 0.0990 M |
| Volume of Bleach sample | 2.00 mL |
| Final Volume of $Na_2S_2O_3$ | 32.62 mL |
| Initial Volume of $Na_2S_2O_3$ | 1.35 mL |
| Volume of $Na_2S_2O_3$ added | |

- A) What is the molarity of NaOCl in this bleach sample?

- B) The density of the bleach solution is 1.10 g/mL. Determine the mass % of NaOCl in the bleach solution.

- C) The manufacturer reports that the bleach is 6.00 % by mass NaOCl. What is the % difference with the manufacturer's claim?

3. (15 points) A series of iron-salicylate solutions were prepared (as we did in the lab) with varying concentrations, and their absorbances measured at 530 nm. The [Fe-SA] (in M) vs. ABS data were plotted using MS Excel and the equation for the line was determined by linear regression. The formula returned by Excel was:

$$y = 1247.5 x$$

A commercial aspirin tablet was then prepared for analysis (as in the experiment that we performed). The tablet was first crushed. The powder was then washed into a 250-mL volumetric flask and excess NaOH solution was added to hydrolyze the acetylsalicylic acid (ASA) to salicylic acid (SA). The volumetric flask was then filled to the mark with DI water. This was called the “stock” solution.

Then, 5.00 mL of the stock solution was pipetted into a 100-mL volumetric flask, and the flask was filled to the mark with an iron (III) chloride solution, converting the SA to the iron-salicylate complex (Fe-SA). This was called the “sample” solution.

The absorbance of the sample solution was then measured at 530 nm, and found to be

$$ABS = 0.739$$

A) What is the [Fe-SA] (in M) in the sample solution?

B) What is the experimental mass of ASA (in mg) in the commercial tablet? The molar mass of ASA ($C_9H_8O_4$) is 180.16 g/mol.


4. (15 points) The following ΔH values were collected in the lab for various acid-base reactions. All species (except water) are aqueous. HCl is a strong acid. CH_3COOH is a weak acid. NaOH is a strong base.

A) Write the net ionic equation for reactions 1 & 2.


B) Calculate the ΔH for reaction 3.

C) In terms of bonds breaking and forming, briefly explain the sign (+ or -) and the magnitude (size) of the three ΔH values.

Explanation:



Net ionic equation:



Net ionic equation:



5. (8 points) You must mix 10 mL of concentrated sulfuric acid (12 *M*) and water to form 0.50 L of a dilute solution.

A) Explain how to do this SAFELY (consider the appropriate lab safety rules in your explanation.)

B) What would be the concentration of the dilute solution?

6. (15 points) In order to determine the molar mass of an unknown volatile liquid, 5.0 mL of the liquid is placed in a pre-massed 125-mL flask capped with aluminum foil with a few small holes punched in it. The system was heated inside a water bath that reached the constant boiling temperature recorded below. The flask with the volatile liquid was removed shortly after it appeared that all of the liquid had vaporized. It was cooled and the flask massed again. The flask was then emptied and its volume determined. Atmospheric pressure in the room was recorded. Based on the data below, calculate the molar mass of the compound.

| | |
|---|------------|
| Mass of flask + aluminum foil | 87.323 g |
| Mass of flask + aluminum foil + condensed vapor | 87.588 g |
| Temperature of boiling water bath | 99.2 °C |
| Volume of the flask | 152 mL |
| Atmospheric pressure | 745.7 mmHg |


7. (2 points) Name several reasons why you move your backpack and belongings away from the work space to the cubbyholes in the back of the labs?

8. (12 points) Oxygen gas is generated by the decomposition of hydrogen peroxide, and collected in a eudiometer tube over water.

The volume of gas inside the tube is 29.8 mL. The height difference between the water in the beaker and in the eudiometer is 33.5 cm (h in the diagram). Atmospheric pressure when this experiment was performed was 735.8 mmHg and the temperature was 22.0°C.

Note: Water vapor pressures at various temperatures are summarized on the constants sheet.

- A) Calculate the total gas pressure inside the tube.



- B) Calculate the pressure of oxygen gas in the tube.